



## Gaviscon Infant

### Uses

Helps to prevent reflux in infants and young children. Indicated for the treatment of gastric regurgitation, gastro-oesophageal reflux and reflux associated with hiatus hernia.

### Dosage

For oral use after mixing with water or milk feed. Infants under 10lbs/4.5kg: 1 sachet. Over 10lbs/4.5kg: 2 sachets. Bottle-fed infants: mix each sachet into 115ml (4 fl. oz.) of feed in the bottle. Shake well. Breast fed infants and other infants up to 2 years: mix each sachet with 5ml (1 teaspoon) of cooled boiled water until a smooth paste is formed, then add another 10ml (2 teaspoons) of cooled boiled water and mix. For breast fed infants, give part way through each feed or meal using a spoon or feeding bottle. For all other infants, give at the end of each meal, using a spoon or feeding bottle. Do not give more than 6 times in 24 hours. Not to be used in premature infants or infants under 1 year except under medical supervision. Not suitable for children over 2 years, adults or the elderly.

### Format and Ingredients

Sachet of powder containing:  
Sodium Alginate 225mg, Magnesium Alginate 87.5mg,

**Manufacturer:**  
RECKITT

**Pack size:** 30 x 2g

**RRP:** £9.83

**Legal Status:** (P)

A division of  
 GROUPE  
EUROCOM

